

The Likely Cause of the EGRET GeV Anomaly and its Implications

F.W. Stecker

NASA Goddard Space Flight Center, Greenbelt, MD 20771

S. D. Hunter

NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

D. A. Kniffen

NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

University Space Research Association, Columbia, MD 21044, USA

Abstract

Analysis of data from the EGRET γ -ray detector on the Compton Gamma Ray Observatory indicated an anomaly in the form of an excess diffuse galactic flux at GeV energies over that which was theoretically predicted. Various explanations for this anomaly have been put forth, including the invocation of supersymmetric dark matter annihilation. We reexamine these explanations here, including a new discussion of the possible systematic errors in the sensitivity determination of the EGRET detector. We conclude that the most likely explanation of the EGRET “GeV anomaly” was an error in the estimation of the EGRET sensitivity at energies above ~ 1 GeV. We give reasons why such a situation could have occurred. We find evidence from our new all-sky analysis which is inconsistent with the assumption that the anomaly can be a signal of supersymmetric dark matter annihilation. We also reconfirm the original results of the EGRET team on the extragalactic γ -ray background spectrum. There are important implications of our analysis for the upcoming Gamma Ray Large Area Telescope (GLAST) mission.

Key words: γ -rays , background radiation

1 Introduction

The EGRET γ -ray detector, a spark chamber telescope flown aboard the Compton Gamma Ray Observatory satellite, provided detailed measurements

of astrophysical γ -ray sources and galactic and extragalactic diffuse γ -ray fluxes. The reported galactic diffuse flux was measured and mapped over the whole sky.

Theoretical studies of the physics and astrophysics of galactic γ -ray production provided predictions of the expected fluxes and energy spectra [1]. In the energy range above ~ 1 GeV, the fluxes reported by the EGRET team were up to $\sim 60\%$ higher than the theoretical predictions [2]. This apparent discrepancy between the reported fluxes and the theoretical calculations will be referred to here as the “GeV anomaly”.

Unresolved galactic point sources can be ruled out as an explanation for the GeV anomaly for two reasons: (1) they would be strongly concentrated in the galactic plane and the GE anomaly, as we will show, is seen isotropically over the whole sky, and (2) the largest class of galactic point sources are pulsars and such sources are concentrated in the inner galaxy and they make up less than 15% of the total galactic flux [3].

There have been three approaches for accounting for the GeV anomaly as a diffuse phenomenon, *viz.* either (1) invoking a cosmic-ray electron source spectrum proportional to E^{-2} with a resulting significant Compton γ -ray component above 1 GeV increasing the flux [4], (2) making modifications to the assumptions regarding both the primary cosmic-ray nucleon and electron spectra in numerical models in order to push up the total theoretical γ -ray fluxes [5], or (3) postulating new physics, namely supersymmetric dark matter annihilation, to account for the anomaly [6].

The first approach postulates that the cosmic-ray electron spectrum observed at Earth is much steeper than the average spectrum in the Galaxy, but that this *ad hoc* situation can result as an effect of the distribution of the supernova remnants (SNR) which produce the electrons and electron propagation effects. A prediction of this model is a center-anticenter asymmetry in the anomaly owing to the strong galactocentric radial distribution of SNR. A reduced bump at high galactic latitudes would also be expected, owing to the steeper local cosmic-ray electron spectrum.

Approaches (2) and (3) have significant implications. Increasing the galactic γ -ray production rate at GeV energies in the model of Ref. [5] produced a reduction in the implied extragalactic diffuse flux in the followup calculation [7], with a marked dip at energies near ~ 1 GeV. This is opposed to the original determination of the extragalactic background spectrum by the EGRET team [8]. Postulating that the GeV anomaly is caused by a γ -ray component from supersymmetric dark matter annihilation has, of course, much greater implications. It is therefore important to reexamine the issue of the GeV anomaly by paying careful attention to the collateral implications of deviations from the

canonical predictions [1] and to examine the more prosaic possibility that the assumed high-energy sensitivity of the EGRET detector may not have been correct. In this paper, we will make the case that a problem in the analysis of the EGRET sensitivity calibration is the most likely explanation for the “GeV anomaly” and that the dark matter hypothesis can be ruled out by an examination of the GeV spectrum over the whole sky.

2 Production of Diffuse Galactic γ -rays

Galactic γ -rays are produced by interactions of relativistic electrons and protons with interstellar gas and photons. The physical processes involved are electron bremsstrahlung, neutral pion production, and Compton interactions of cosmic ray electrons [9],[10],[1]. Of these processes, the decay of neutral pions produced by cosmic rays interacting with interstellar gas is expected to dominate at energies above 0.1 GeV [1],[2]. The vast majority of these γ -rays are produced by cosmic rays with energies below ~ 20 GeV [11]. In this energy range the cosmic ray spectrum, particularly in the local galactic neighborhood that accounts for the γ -ray production at high galactic latitudes, is well measured [12]. The pion production cross section at these energies is also very well known [11],[13]. At γ -ray energies up to ~ 1 GeV, the pion production process is well described by the “isobar-plus-fireball” model [9],[14],[15]. At energies above ~ 1 GeV, it is generally assumed that scaling holds and that the γ -ray spectrum will have the same spectral index as the primary proton spectrum [15],[16]. More recent calculations [18] are in good agreement with the results presented in Refs. [15] and [16], confirming that the physics of pion production at GeV energies is well determined. A modification of this physics involving scaling violation has been suggested [17], however the resulting effect is too small to explain the GeV anomaly by scaling violation alone.

It has been suggested that the GeV anomaly can be explained by postulating that the average primary proton spectrum in the galaxy is harder than that observed locally. With scaling assumed, the required proton spectral index is $\Gamma = 2.45$ [18],[19]. Even with some scaling violation [17], one requires a spectral index $\Gamma \simeq 2.5$. The problem with this assumption is that the local proton spectrum has a significantly steeper measured index $\Gamma = 2.76 \pm 0.03$ [12]. At galactic latitudes above $\sim 20^\circ$ from the plane, the diffuse γ -rays are produced by these locally measured protons (and a smaller number of cosmic-ray α particles which have a similar spectral index [12]). Even should the proton spectrum in the inner galaxy be harder than the local spectrum at high galactic latitudes, it is the steeper local spectrum which produces the high-latitude pion-decay γ -rays. Therefore, the predicted spectrum should be the canonical one. In other words, there should be no GeV anomaly at high galactic latitudes. This conclusion would also result if the anomaly is strictly

produced by a harder cosmic-ray electron spectrum averaged over the whole galactic disk [4].

On the contrary, it is generally accepted that the GeV anomaly is *uniform over the entire sky*, a result which was obtained from the EGRET data and which was implicit in Figs. 3 and 5 of Ref. [8] by the EGRET group. We will show here, in a very quantitative way, that the anomaly is seen at *all* galactic latitudes independent of the line-of-sight density of galactic hydrogen gas. We will further argue that such complete isotropy is what would be expected if this anomaly is most likely traceable to the detector itself.

3 The Celestial Distribution of the GeV Anomaly

Our detailed analysis of the galactic γ -ray flux has now been expanded to cover the entire sky [22]. This was accomplished by extending the latitude range of the galactic plane analysis to $\pm 25^\circ$, chosen to match the latitude range which includes all significant emission by molecular hydrogen clouds [21], and further extending the model to the galactic poles [22] by using the all-sky Leiden-Dwingaloo [23] and Instituto Argentino de Radioastronomia (IAR) Southern Hemisphere HI survey [24]. The three dimensional cosmic-ray density for the latitude range $|b| < 25^\circ$ was derived from the Galactic plane matter distribution on the assumption of dynamic balance [25]. The γ -ray production function per H atom in the $22^\circ < |b| < 25^\circ$ latitude range was extended to the poles on the assumption that γ -ray production in this range is dominated by interactions with cosmic rays having the locally measured spectrum. The analysis approach used in Ref. [8] to determine the extragalactic diffuse emission was then repeated using the new all-sky model for each of the ten standard energy ranges as well as the four broad energy ranges discussed in Ref. [2]. This analysis, using the EGRET phase 1 and phase 2 source-subtracted data, yields an extragalactic γ -ray spectrum which is consistent with the one derived in Ref. [8]. This consistency is expected because the extragalactic flux is the residual observed flux as the galactic model flux is extrapolated to zero. The slope of the extrapolation does not affect the intercept, as shown in Figure 1. Furthermore, this all-sky analysis confirms that the GeV anomaly is *uniform over the entire sky*, a result which was implicit in Figs. 3 and 5 Ref. [8], but which is now more quantitatively shown in Fig. 1. The lack of any structure in the anomaly related to the Galactic plane, galactic center, anti-center or halo, strongly indicates that the GeV anomaly is due to a systematic error in the EGRET calibration (see below) rather than being a real astronomical effect. The excellent 1:1 correlation of the EGRET data with the expanded all-sky model over the entire sky and the entire range of emission, after multiplication by a single renormalization factor, further corroborates this conclusion as also shown in Fig. 1. Table 1 and Fig. 2 breaks down this effect as a func-

tion of energy which indicates a miscalibration which gets worse at the higher energies.

The overall effect of applying this renormalization correction for point sources is slight over a large energy range on a log-log plot and does not conflict with any theoretical models within the large observational error bars of the data above 1 GeV. For γ -ray blazars observed by EGRET, the fit to power-law spectra would result in slightly steeper spectra, with the error bars on specific points generally large in the energy range above 1 GeV.

To specifically illustrate the effect of our renormalization factors on the spectra of point sources, we have applied our renormalization to the EGRET data on both the pulsed and unpulsed nebula spectrum of the Crab Nebula as shown in Figure 3 based on the compilation in Ref. [26]. The renormalized spectrum is consistent with previous theoretical models of γ -ray emission from the Crab Nebula [26],[27].

4 The Dark Matter Hypothesis

It has been suggested that the GeV anomaly is the result of a new component to the diffuse γ -ray spectrum, one that is in addition to the pion-decay component, namely a component from supersymmetric dark matter annihilation [6]. However, there are serious problems with this hypothesis: (1) The same process of dark matter annihilation would produce a flux of cosmic-ray antiprotons which is incompatible with the measured value [28]. (2) The celestial γ -ray distribution from dark matter annihilation would be both highly asymmetric and clumped [29]. This is again in direct contradiction with isotropy of the anomaly.

5 The Energy Calibration of the EGRET Detector

It was pointed out in the EGRET calibration paper [30] that at high energies the EGRET sensitivity is poorly known, partly due to the uncertain effect of self-vetoing caused by charged-particle backplash impinging onto the anticoincidence scintillator following shower generation in the detector [8]. This effect eliminates good γ -ray events. Any uncertainty in the self-veto correction would alter the sensitivity of the detector at energies above ~ 1 GeV.

An attempt to quantify the amount of backplash was performed experimentally up to 10 GeV during the calibration of EGRET at the Stanford Linear

Accelerator Center (SLAC) with over a thousand hours of γ -ray beam exposure. The beam was scanned across the face of the tracking detector at a series of off-axis angles from 0° to 40° with three different roll-axis angles and 10 different energies. An EGS (Electron Gamma Shower) Monte Carlo (MC) code was used to study the effect of the backscatter under differing conditions, but the fidelity of the MC model was low. It did indicate that the backscatter was strongly dependent on the charged-particle threshold, which was known to be strongly variable across the anti-coincidence detector. The photomultiplier gains were reduced after the SLAC calibration to mitigate the higher than anticipated backscatter effect. The effect of lowering the gains is substantial, and is not reflected in the sensitivity calibration results as given in Ref. [30].

The all-sky pervasiveness of the apparent excess flux above 1 GeV energy (see above) strongly indicates that this anomaly is intrinsic to the detector. A systematic error in the understanding of the detector response would be one likely source. In an effort to investigate this possibility, we have reexamined the EGRET archival data carefully, selecting only the highest quality EGRET data. The selection was based on: (1) choosing γ -ray events with the most reliable energy determination, (2) selecting γ -ray events within 30° of the detector axis for which the instrument response is most reliably determined, and (3) excluding γ -ray events within a 4σ angular uncertainty from the Earth's horizon. We have examined data on the pulsed and unpulsed emission from the Crab, Vela, and Geminga and the diffuse flux from the Lockman Hole for phases 1 – 4 of the EGRET data for energies $> 1\text{GeV}$. We find that for different observations of these sources, there were variations in their measured fluxes in excess of 40%. Since the flux from these sources is expected to be constant, this indicates systematic errors caused apparent time variations over different observing periods. This is not surprising, since the effect of degraded performance with the aging of the spark chamber gas was calibrated in flight by fixing the > 100 MeV flux and not taking account of the energy dependence of this effect [31], thus producing a false variability at GeV energies.

While we find anomalous behavior pointing to an imprecise knowledge of the detector response, we are unable to explain the GeV anomaly as caused by any single systematic instrument effect that we have studied. Other contributors besides an imprecise knowledge of the detector response that could explain an apparent trans-GeV anomaly include (1) an error in the analysis which might cause non- γ -ray events to be identified as good high-energy events, (2) an incorrect assignment of energies and (3) an induced detector background in the harsh environment of a low-Earth orbit. It is difficult to eliminate any of these possibilities, but no specific evidence has been found that they exist.

Given the systematic energy dependent uncertainties in the sensitivity calibration for the EGRET instrument discussed in this paper, the published EGRET results above 1 GeV cannot be reliably depended upon. We suggest

Table 1
 INVERSE NORMALIZATION FACTOR VS.ENERGY

Energy Bin	Inverse Normalization Factor
30-50 MeV	0.815 ± 0.009
50-70 MeV	0.992 ± 0.018
70-100 MeV	0.947 ± 0.019
100-150 MeV	0.959 ± 0.019
150-300 MeV	0.925 ± 0.013
300-500 MeV	0.887 ± 0.022
500-1000 MeV	0.965 ± 0.026
1-2 GeV	1.26 ± 0.06
2-4 GeV	1.57 ± 0.17
4-10 GeV	1.85 ± 0.46

that one should incorporate the energy dependent correction factor shown in Figure 2. This suggestion applies to possible use by the GLAST LAT (Large Area Telescope) collaboration for comparison as a sensitivity calibration check on their extensive Monte Carlo simulations and beam tests. We must await results from GLAST in order to pursue this question further.

6 Conclusions

Our new all-sky analysis of the EGRET data confirms a systematic anomaly in the form of an apparent excess galactic emission at energies above 1 GeV over what is expected from galactic cosmic ray interactions with gas nuclei in atomic and molecular clouds and interstellar photons, as previously reported [2]. We find further quantitative support that this anomaly is constant over the whole sky, not being correlated with any astronomical or galactic features. It is therefore most likely caused by a systematic error in the calibration of the effective detector sensitivity. Although a detailed reanalysis of the calibration data is impossible at this time, we have shown that plausible systematic uncertainties in the calibration of the EGRET sensitivity for γ -ray energies above 1 GeV can readily account for the universal anomalous excess flux. We conclude that neither making modifications to the observed primary cosmic-ray nucleon and electron spectra in order to push up the predicted theoretical γ -ray flux [5] nor postulating a component from supersymmetric dark matter annihilation [6] are required to account for the “GeV anomaly”.

Our new analysis is consistent with the extragalactic flux derived in Ref. [8]. This reevaluation of the EGRET sensitivity also has implications for calibration checks of the GLAST γ -ray detector to be launched in the near future.

Acknowledgments

We thank Michael Salamon and Dave Thompson for their comments and suggestions regarding this work and Zev Gurman for his analysis of EGRET archival data.

References

- [1] F. W. Stecker, *Astrophys. J.* **212**, 60 (1977).
- [2] S. D. Hunter *et al.*, *Astrophys. J.* **481**, 205 (1997).
- [3] A. K. Harding and F. W. Stecker, *Nature* **290**, 316 (1981).
- [4] M. Pohl and J. A. Esposito, *Astrophys. J.* **507**, 327 (1998).
- [5] A. W. Strong, I. V. Moskalenko, and O. Reimer, *Astrophys. J.* **613**, 962 (2004).
- [6] W. De Boer, *Astron. and Astrophys.* **444**, 51 (2005).
- [7] A. W. Strong, I. V. Moskalenko, and O. Reimer, *Astrophys. J.* **613**, 956 (2004).
- [8] P. Sreekumar *et al.*, *Astrophys. J.* **494**, 523 (1998).
- [9] F. W. Stecker, *Astrophys. and Space Sci.* **6**, 377 (1970).
- [10] F. W. Stecker, *Cosmic Gamma Rays* (Baltimore: Mono Pub.) (1971).
- [11] F. W. Stecker, *Astrophys. J.* **185**, 499 (1973).
- [12] Y. Shikaze *et al.*, *Astropart. Phys.* **28**, 154 (2007).
- [13] V. de Ataíde Malafaia Lopes dos Santos, e-print [astro-ph/nuc1-th/0602026](#).
- [14] C. D. Dermer, *Astron. and Astrophys.* 157 (1986) 223.
- [15] F. W. Stecker, in *Cosmic Gamma Rays, Neutrinos, and Related Astrophysics* ed. M. M. Shapiro and J. P. Wefel (Dordrecht: Kluwer), 85 (1989).
- [16] F. W. Stecker, *Astrophys. J.* **228**, 919 (1979).
- [17] T. Kamae *et al.*, *Astrophys. J.* **647**, 692 (2006).
- [18] M. Mori, *Astrophys. J.* **478**, 225 (1997).

- [19] P. Gralewicz, J. Wdowczyk, A. W. Wolfendale and L. Zhang, *Astron. and Astrophys.* **318**, 925 (1997).
- [20] In Ref.[5] the approach is taken of arbitrarily raising the cosmic-ray proton and electron spectra by different amounts in order to optimize the fit to the reported EGRET GeV anomaly. We see no basis for these *ad hoc* assumptions, particularly as the proton flux is well measured [12].
- [21] T. M. Dame *et al.*, *Astrophys. J.* **322**, 706 (1987).
- [22] S. D. Hunter, D. A. Kniffen and F. W. Stecker, in preparation.
- [23] D. Hartmann and B. W. Burton 1997, *Atlas of Galactic Neutral Hydrogen*, Cambridge, UK: Cambridge U. Press
- [24] R. Morras, E. Bajaja, E. M. Arnal, and W. G. L. Pöppel, *Astron. and Astrophys. Sup.*, **142**, 25 (2000).
- [25] D. L. Bertsch *et al.*, *Astrophys. J.* **416**, 587, (1993).
- [26] L. Kuiper *et al.*, *Astron. and Astrophys.* **378**, 918 (2001).
- [27] O. C. de Jager *et al.*, *Astrophys. J.* **457**, 253 (1996).
- [28] L. Bergström, J. Edsjö, M. Gustafsson and P. Salati, *Jour. Cosm. and Astropart. Phys.* 05(2006)006.
- [29] J. Diemand, M. Kuhlen, and P. Madau, *Astrophys. J.* **657**, 262 (2007).
- [30] D. J. Thompson *et al.* *Astrophys. J. Suppl.* **86**, 629 (1993).
- [31] J. A. Esposito *et al.* *Astrophys. J. Suppl.* **123**, 203 (1999).
- [32] Y. C. Lin, EGRET internal document, unpublished (1992).




Fig. 1. (a) Plot of integral ($E > 1$ GeV), all-sky diffuse model flux vs. EGRET observed flux for $335^\circ < l < 45^\circ$, $|b| < 90^\circ$. (b) A similar plot with a renormalization factor of $(1.6)^{-1}$ applied to the observed flux. In both plots, the integral extragalactic diffuse flux of $1.5 \times 10^{-6} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ has been added to the diffuse model. The dotted line indicates the expected 1:1 relationship between the model and observed fluxes. Contours show the number of $0.5^\circ \times 0.5^\circ$ pixels containing the flux indicated within a bin of width $1.6 \times 10^{-6} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$. The contour values are $10^4, 10^3, 10^2, 10^1, \dots$. These plots clearly demonstrate that the GeV anomaly exists uniformly over the whole sky and extends from high to low intensity galactic flux emission.




Fig. 2. Required inverse renormalization factor for different energy bins, given as the ratio of observed-to-predicted flux *vs.* energy.




Fig. 3. Pulsed and Nebula spectrum of the Crab Nebula with and without the correction factors from Figure 2 applied to the EGRET data, based on the compilation in Ref. [26]. The crosses in the energy range from 50 MeV to 10 GeV show the original EGRET data points and the diamonds show the renormalized values.